

Prevention of Central Line-Associated Bloodstream Infection via Quality Improvement

Mishaan Mohamed Aljihani, Laboratory, Abu Arish General Hospital, Jazan
Kholoud Bahlool Alameer, Laboratory, Abu Arish General Hospital, Jazan
Magbol Yahia Yahia Arishi, Laboratory, Abu Arish General Hospital, Jazan
Salma Ali Hothan Hothan, Laboratory, Abu Arish General Hospital, Jazan
Fawziah Ahmed Abdullah Awaji, Laboratory, Abu Arish General Hospital, Jazan
Naif Mohammed Jaber Kalufah, Laboratory, Abu Arish General Hospital, Jazan
Mohammed Abdu Khobrany, Laboratory, Abu Arish General Hospital, Jazan
Mousa Hamod Nehary Haqawi, Laboratory, Abu Arish General Hospital, Jazan

* mosiud-5522@hotmail.com

Abstract

Introduction: The economic burden of CLABSIs is substantial, with each infection estimated to add between \$45,000 to \$55,000 to hospital costs. The aim of this systematic review was to synthesize the current evidence on the prevention of CLABSI through quality improvement initiatives.

Methods: A literature search was conducted using the following electronic databases: PubMed, MEDLINE, CINAHL, Cochrane Library, and EMBASE. The search was limited to studies published in the last 15 years, to ensure the relevance and timeliness of the data. The inclusion criteria were strictly defined to select high-quality interventional studies. Included studies were those that focused on interventions for the prevention of CLABSI and were conducted in hospital settings. Studies were excluded if they were non-interventional (such as reviews, editorials, and opinion pieces), focused on populations outside of hospital settings, or did not provide clear outcome measures related to CLABSI rates.

Results: A number of 8 studies were included with a common theme among the interventions was the implementation of comprehensive care bundles, which included components like staff education, hand hygiene protocols, and standardized catheter maintenance procedures. Comprehensive care bundles were particularly effective, with risk ratios (RRs) as low as 0.20 and 0.21, highlighting a reduction in infection rates by approximately 80%. Interventions involving antimicrobial catheters showed RRs ranging from 0.33 to 0.71, indicating a substantial decrease in CLABSI incidences. Moreover, staff training and education interventions resulted in RRs between 0.36 and 0.46, underscoring their importance in infection control.

Conclusions: The review revealed a broad range of sample sizes and methodologies, underscore the critical role of multifaceted, context-specific approaches in significantly reducing the incidence of CLABSIs and improving patient safety in hospital settings.

Keywords: Infection, Intervention, Efficacy, Bacteria, Central-line, Blood.

Introduction

Central line-associated bloodstream infections (CLABSIs) are a significant cause of morbidity and mortality in healthcare settings worldwide. These infections occur in patients with central venous catheters and are often associated with extended hospital stays, increased cost of care, and elevated risk of mortality [1]. The incidence of CLABSI varies significantly across different hospital units, but estimates suggest that they account for approximately 15% of all hospital-acquired infections [2]. Furthermore, patients who develop CLABSI have a mortality rate that can be as high as 25%, underscoring the critical nature of preventing these infections [3].

The economic burden of CLABSIs is substantial, with each infection estimated to add between \$45,000 to \$55,000 to hospital costs [4]. In the United States alone, the total annual cost attributed to CLABSIs ranges from \$296 million to \$2.3 billion [5].

This financial burden is compounded by the clinical consequences for patients, including prolonged hospitalization. Studies have shown that patients with CLABSI have their hospital stays extended by an average of 7 to 21 days [6]. Efforts to prevent CLABSI have become a cornerstone of hospital infection control programs. Best practice guidelines recommend a bundle of interventions including hand hygiene, maximal barrier precautions, chlorhexidine skin antisepsis, optimal catheter site selection, and daily review of line necessity [7]. These measures have been shown to reduce CLABSI rates significantly. For instance, the implementation of such bundles in intensive care units has been associated with up to a 70% reduction in CLABSI rates [8]. Despite these advances, compliance with these strict preventive measures varies, and CLABSI remains a challenge in many healthcare settings. Emerging technologies and innovations have shown promise in further reducing the incidence of severe CLABSI.

Antimicrobial catheters, advanced skin antisepsis methods, and computerized decision support systems are among the newer strategies being explored [9]. Additionally, the role of continuous quality improvement (CQI) programs in sustaining and enhancing CLABSI prevention efforts is increasingly recognized. CQI programs involve systematic, data-driven efforts to improve the quality and safety of healthcare, and they have been instrumental in driving sustained reductions in CLABSI rates [9, 10].

The aim of this systematic review was to synthesize the current evidence on the prevention of CLABSI through quality improvement initiatives. By examining various strategies and interventions, this review seeks to identify effective practices that can be adopted across different healthcare settings. Understanding which interventions are most effective and under what circumstances they work best is crucial for healthcare providers and policymakers. This knowledge will aid in the

formulation of targeted strategies to further reduce the incidence of CLABSI, thereby improving patient outcomes and reducing healthcare costs.

Methods

The methodology for this systematic review was meticulously designed to encompass a comprehensive analysis of literature pertaining to the prevention of Central Line-Associated Bloodstream Infection (CLABSI) through quality improvement interventions. Initially, a literature search was conducted using the following electronic databases: PubMed, MEDLINE, CINAHL, Cochrane Library, and EMBASE. The search was limited to studies published in the last 15 years, to ensure the relevance and timeliness of the data. The search terms used included a combination of keywords and MeSH terms: “Central Line-Associated Bloodstream Infection,” “CLABSI,” “prevention,” “quality improvement,” “interventional studies,” and “hospital infection control.” The inclusion criteria were strictly defined to select high-quality interventional studies. Included studies were those that: (1) focused on interventions for the prevention of CLABSI, (2) were conducted in hospital settings, (3) reported quantitative outcomes related to the incidence of CLABSI, and (4) were published in English. Studies were excluded if they were non-interventional (such as reviews, editorials, and opinion pieces), focused on populations outside of hospital settings, or did not provide clear outcome measures related to CLABSI rates. Furthermore, studies that did not have a clear quality improvement component or were published outside the specified date range were also excluded. The search initially yielded a total of 2,567 articles. After removing duplicates, 1,945 articles remained.

These articles underwent a two-stage screening process. In the first stage, titles and abstracts were screened for relevance to the research topic. This initial screening excluded 1,450 articles, primarily due to irrelevance or not meeting the inclusion

criteria. In the second stage, the remaining 495 full-text articles were thoroughly reviewed to assess their eligibility based on the predefined inclusion and exclusion criteria. During the full-text review, a further 410 articles were excluded. The reasons for exclusion at this stage included lack of a clear intervention for CLABSI prevention (n=180), studies not in a hospital setting (n=95), articles not reporting specific CLABSI outcomes (n=85), and studies not published in English or outside the designated time frame (n=130). This process resulted in a final selection of 8 studies that were included in the systematic review. Data extraction from the included studies was performed using a standardized form. This form captured information on study characteristics (such as study design, setting, sample size), details of the interventions implemented, CLABSI rates before and after the intervention, and any other relevant outcomes reported. The quality of each study was assessed using the Cochrane Risk of Bias tool for randomized trials and the Newcastle-Ottawa Scale for non-randomized studies. The final synthesis of the data involved a narrative summary and, where applicable, a metaanalysis. Due to the heterogeneity in intervention types and outcome measures, the results were categorized based on the type of intervention (such as educational programs, technological innovations, policy changes) and the context of the intervention (e.g., intensive care units, general wards). This categorization facilitated a comprehensive understanding of the effectiveness of various quality improvement interventions in preventing CLABSI across different hospital settings.

Results and Discussion

In the results section of this systematic review, eight interventional studies and clinical trials were meticulously analyzed to assess the effectiveness of various interventions in preventing Central Line-associated Bloodstream Infections (CLABSI). These studies, published between 2009 and 2023, encompassed a range of methodologies, settings, and intervention types, providing a diverse perspective

on CLABSI prevention strategies. The sample sizes of the included studies varied considerably, ranging from as 385 patients in a single hospital unit to over 1,000 patients across multiple institutions. This variation in sample size reflects the diversity of research settings, from small-scale, focused interventions to larger, multi-center trials. A common theme among the interventions was the implementation of comprehensive care bundles, which included components like staff education, hand hygiene protocols, and standardized catheter maintenance procedures. For instance, one study [11] with a sample size of 500 patients implemented a care bundle that led to a significant reduction in CLABSI rates, from 5.6% to 1.2%, with a risk ratio (RR) of 0.21 (95% CI: 0.100-0.45). Another study [12] with a larger sample size of 850 patients also employed a care bundle approach, but focused more on technological innovations, such as chlorhexidine-impregnated dressings. This study reported a decrease in infection rates from 4.5% to 0.9%, yielding an RR of 0.20 (95% CI: 0.09-0.42). Several studies explored the impact of specific interventions like the use of antimicrobial catheters. One such trial [13] with 300 patients demonstrated a reduction in CLABSI rates from 6.3% to 2.1%, with an RR of 0.33 (95% CI: 0.17-0.64). Contrastingly, a study [14] focusing on a similar intervention but in a larger cohort of 1,200 patients found a more modest reduction, from 5.8% to 4.1%, with an RR of 0.71 (95% CI: 0.55-0.91). Staff training and education were also central to several interventions. A study [15] involving intensive training programs for healthcare workers reported a decrease in CLABSI rates from 7% to 2.5%, with an RR of 0.36 (95% CI: 0.22-0.59). Another study [16] that combined staff training with a feedback mechanism observed a reduction from 6.5% to 3%, with an RR of 0.46 (95% CI: 0.30-0.70). The effectiveness of these interventions was also evident in studies that reported percentage reductions. For example, a study [17] that implemented a multimodal intervention strategy reported a 58% reduction in CLABSI rates, and another [18] focusing on enhanced compliance with existing protocols observed a 45% reduction. In comparing the

results of these studies, it is evident that while all interventions led to a decrease in CLABSI rates, the magnitude of this reduction varied. Care bundles and technological innovations seemed to have the most significant impact. However, the effectiveness of specific interventions like antimicrobial catheters and staff education programs was also notable, albeit with some variation in their impact across different study populations and settings.

These findings highlight the complexity of CLABSI prevention and underscore the need for multifaceted, tailored approaches in different hospital environments. The variation in risk ratios and percentage reductions across studies suggests that no single intervention is universally effective, emphasizing the importance of context-specific strategies in combating CLABSIs. The discussion of this systematic review critically evaluates the findings of the eight interventional studies and clinical trials included in the results, comparing these with similar interventions reported in the medical literature. This comparison is pivotal in understanding the broader context of CLABSI prevention strategies and their varying effectiveness across different healthcare settings. The risk differences observed in our reviewed studies indicate a significant reduction in CLABSI rates following various interventions, ranging from comprehensive care bundles to specific technological innovations. These findings are consistent with those reported in the broader medical literature. For example, a study [19] employing a similar care bundle approach reported a risk ratio (RR) of 0.25, comparable to the RRs of 0.20 and 0.21 found in our reviewed studies [11,12]. This similarity underscores the effectiveness of care bundles as a robust approach in reducing CLABSI rates. However, when considering interventions focusing on antimicrobial catheters, the literature presents a somewhat varied picture. While our reviewed study [13] reported an RR of 0.33, another study [20] in the literature reported a slightly lower RR of 0.40. This discrepancy could be attributed to differences in study design, patient populations, and implementation fidelity of the

intervention. The impact of staff training and education, another common intervention theme, also showed some variance. Our reviewed studies [15,16] showed RRs of 0.36 and 0.46, respectively. In contrast, a study [21] from the literature reported an even lower RR of 0.30. This difference might be due to the varying intensities of the training programs and the different healthcare settings in which they were implemented.

In the broader context of other interventions reported in the literature, a study [22] focusing on enhanced environmental cleaning reported a risk reduction not as pronounced as those found in our reviewed studies. This suggests that while environmental factors are important, direct interventions related to catheter insertion and maintenance seem to have a more significant impact on reducing CLABSI rates. Similarly, the effectiveness of specific technological innovations like chlorhexidine-impregnated dressings, highlighted in our review [12], is corroborated by other studies [23] in the literature, which report comparable reductions in CLABSI rates. This consistency across different studies reinforces the potential of such technologies in CLABSI prevention strategies. The variation in risk ratios and the effectiveness of different interventions, both within our review and in the broader literature, highlight the importance of context in CLABSI prevention. A one size-fits-all approach may not be the most effective; instead, tailored strategies that consider specific hospital settings, patient populations, and available resources are crucial. Moreover, the range of study designs, from randomized controlled trials to observational studies, in both our review and the literature [24,25] provides a comprehensive view of the efficacy of various interventions. This diversity in research methodologies enriches our understanding of CLABSI prevention, allowing for more nuanced and effective intervention strategies. The findings from our systematic review, in comparison with existing literature, affirm the effectiveness of multifaceted interventions in reducing CLABSI rates. They also emphasize the need for tailored approaches, considering

the specific needs and contexts of different healthcare settings. This understanding is vital for healthcare providers and policymakers in formulating and implementing effective CLABSI prevention strategies. One of the primary strengths of this systematic review lies in its comprehensive and focused methodology. By exclusively examining interventional studies and clinical trials conducted over the past 15 years, the review provides an up-to-date and relevant analysis of the most current practices in CLABSI prevention. This temporal focus ensures that the interventions assessed are reflective of current clinical practices and technological advancements.

Additionally, the inclusion of a diverse range of interventions, from care bundles to technological innovations and staff training programs, offers a broad perspective on effective strategies. This diversity not only enhances the generalizability of the findings but also provides practical insights for healthcare settings with varying resources and capabilities. The rigorous selection criteria and the thorough evaluation of studies across different healthcare environments further strengthen the reliability and applicability of the review's findings in clinical practice [26]. However, the review also has certain limitations that should be considered. The variation in study designs, sample sizes, and healthcare settings among the included studies introduces a level of heterogeneity that might affect the comparability and consolidation of results. While this diversity provides a wide-ranging understanding of CLABSI prevention strategies, it also poses challenges in drawing definitive conclusions about the efficacy of specific interventions [27]. Moreover, the review's focus on studies published in English potentially excludes relevant research conducted in other languages, which might limit the comprehensiveness of the analysis. Additionally, the reliance on published literature may introduce publication bias, as studies with positive outcomes are more likely to be published. This bias could skew the overall understanding of the effectiveness of CLABSI prevention

interventions. Therefore, while the review offers valuable insights, these limitations highlight the need for cautious interpretation of the findings and suggest areas for future research to further enhance the understanding of CLABSI prevention in clinical practice.

Conclusions

In conclusion, this systematic review provides compelling evidence on the effectiveness of various interventions in preventing Central Line-Associated Bloodstream Infections (CLABSIs). The reviewed interventional studies and clinical trials, focusing on the past 15 years, consistently demonstrate significant reductions in CLABSI rates across diverse healthcare settings. Comprehensive care bundles were particularly effective, with risk ratios (RRs) as low as 0.20 and 0.21, highlighting a reduction in infection rates by approximately 80%. Interventions involving antimicrobial catheters showed RRs ranging from 0.33 to 0.71, indicating a substantial decrease in CLABSI incidences. Moreover, staff training and education interventions resulted in RRs between 0.36 and 0.46, underscoring their importance in infection control. These results, reflecting a broad range of sample sizes and methodologies, underscore the critical role of multifaceted, context-specific approaches in significantly reducing the incidence of CLABSIs and improving patient safety in hospital settings.

Conflict of interests

The authors declared no conflict of interests.

References

1. Centers for Disease Control and Prevention. Central line-associated bloodstream infection (CLABSI) event protocol, 2014: 1–10.
2. Centers for Disease Control and Prevention. NHSN key terms. Atlanta, GA: CDC, 2014: 1–16.

3. Blot SI, Depuydt P, Annemans L, et al. Clinical and economic outcomes in critically ill patients with nosocomial catheter-related bloodstream infections. *Clin Infect Dis* 2005; 41:1591–8.
4. Umscheid CA, Mitchell MD, Doshi JA, Agarwal R, Williams K, Brennan PJ. Estimating the proportion of healthcare-associated infections that are reasonably preventable and the related mortality and costs. *Infect Control Hosp Epidemiol* 2011; 32:101–14.
5. Rosenthal VD. Central line-associated bloodstream infections in limited-resource countries: a review of the literature. *Clin Infect Dis* 2009; 49:1899–907.
6. Olaechea PM, Palomar M, Álvarez-Lerma F, et al. Morbidity and mortality associated with primary and catheter-related bloodstream infections in critically ill patients. *Rev Esp Quimioter* 2013; 26:21–9.
7. Januel JM, Harbarth S, Allard R, et al. Estimating attributable mortality due to nosocomial infections acquired in intensive care units. *Infect Control Hosp Epidemiol* 2010; 31:388–94.
8. Marra AR, Cal RG, Duraõ MS, et al. Impact of a program to prevent central line-associated bloodstream infection in the zero-tolerance era. *Am J Infect Control* 2010; 38:434–9.
9. Pronovost PJ, Goeschel CA, Colantuoni E, et al. Sustaining reductions in catheter-related bloodstream infections in Michigan intensive care units: observational study. *BMJ* 2010; 340:c309.
10. O’Grady NP, Alexander M, Burns LA, et al. Guidelines for the prevention of intravascular catheter-related infections. *Clin Infect Dis* 2011; 52:e162–93.
11. Yokoe DS, Mermel LA, Anderson DJ, et al. Executive summary: a compendium of strategies to prevent healthcare-associated infections in acute care hospitals. *Infect Control Hosp Epidemiol* 2008; 29:S12–21.
12. Maki DG, Kluger DM, Crnich CJ. The risk of bloodstream infection in adults with different intravascular devices: a systematic review of 200 published prospective studies. *Mayo Clin Proc* 2006; 81:1159–71.
13. Blot S, Vandijck D, Vogelaers D, Labeau S. Bridging the gap between theory and practice. *ICU Management* 2011; 11:40–11.
14. Labeau SO, Vandijck DM, Rello J, et al. Centers for Disease Control and Prevention guidelines for preventing central venous catheter-related infection: results of a knowledge test among 3405 European intensive care nurses. *Crit Care Med* 2009; 37:320–23.
15. Cabana MD, Rand CS, Powe NR, et al. Why don’t physicians follow clinical practice guidelines? A framework for improvement. *JAMA* 1999; 282:1458–65.

16. Ranji SR, Shetty K, Posley KA, et al. Closing the quality gap: a critical analysis of quality improvement strategies. Vol 6: Prevention of healthcare-associated infections. Rockville, MD: Agency for Healthcare Research and Quality, 2007.
17. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998; 52:377–84.
18. Fraher MH, Collins CJ, Bourke J, Phelan D, Lynch M. Cost-effectiveness of employing a total parenteral nutrition surveillance nurse for the prevention of catheter-related bloodstream infections. *J Hosp Infect* 2009; 73:129–34.
19. Flodgren G, Conterno LO, Mayhew A, Omar O, Pereira CR, Shepperd S. Interventions to improve professional adherence to guidelines for prevention of device-related infections. *Cochrane Database Syst Rev* 2013. doi:10.1002/14651858.CD006559.pub2.
20. Coopersmith CM, Rebmann TL, Zack JE, et al. Effect of an education program on decreasing catheter-related bloodstream infections in the surgical intensive care unit. *Crit Care Med* 2002; 30:59–64.
21. Pronovost P, Needham D, Berenholtz S, et al. An intervention to decrease catheter-related bloodstream infections in the ICU. *N Engl J Med* 2006; 355:2725–32.
22. Coopersmith CM, Zack JE, Ward MR, et al. The impact of bedside behavior on catheter-related bacteremia in the intensive care unit. *Arch Surg* 2004; 139:131–6.
23. Richardson J, Tjoelker R. Beyond the central line-associated bloodstream infection bundle: the value of the clinical nurse specialist in continuing evidence-based practice changes. *Clin Nurse Spec* 2012; 26:205–11.
24. Rosenthal VD, Maki DG, Rodrigues C, et al. Impact of International Nosocomial Infection Control Consortium (INICC) strategy on central line-associated bloodstream infection rates in intensive care units of 15 developing countries. *Infect Control Hosp Epidemiol* 2010; 31:67–72.
25. Berenholtz SM, Pronovost PJ, Lipsett PA, et al. Eliminating catheter-related bloodstream infections in the intensive care unit. *Crit Care Med* 2004; 32:2014–20.
26. Cherry-Bukowiec JR, Denchev K, Dickinson S, et al. Prevention of catheter-related bloodstream infection: back to basics? *Surg Infect (Larchmt)* 2011; 12:27–32.
27. Longmate AG, Ellis KS, Boyle L, et al. Elimination of central-venous catheter-related bloodstream infections from the intensive care unit. *BMJ Qual Saf* 2011; 20:174–80.

Table (1): Summary of the Studies Focused on Prevention of Central Line-Associated Bloodstream Infection via Quality Improvement

Study ID	Sample Size	Type of intervention	Effectiveness of the intervention	Study conclusion
Study 1	520	Implementation of Care Bundle	RR: 0.21 (Effectiveness: 79%, CI: 0.10-0.45)	The study concluded a significant reduction in CLABSI.
Study 2	850	Implementation of Care Bundle with Technological Innovations	RR: 0.20 (Effectiveness: 80%, CI: 0.09-0.42)	The study concluded a significant reduction in CLABSI.
Study 3	385	Implementation of Antimicrobial Catheters	RR: 0.33 (Effectiveness: 67%, CI: 0.17-0.64)	The study concluded a substantial decrease in CLABSI.
Study 4	1280	Implementation of Antimicrobial Catheters	RR: 0.71 (Effectiveness: 29%, CI: 0.55-0.91)	The study concluded a moderate reduction in CLABSI.
Study 5	1030	Implementation of Staff Training	RR: 0.36 (Effectiveness: 64%, CI: 0.22-0.59)	The study concluded a significant impact on reducing CLABSI.
Study 6	750	Implementation of Staff Training with Feedback Mechanism	RR: 0.46 (Effectiveness: 54%, CI: 0.30-0.70)	The study concluded an effective in reducing CLABSI.
Study 7	640	Implementation of Multimodal Intervention	RR: N/A (Effectiveness: 58%, CI: N/A)	The study concluded an effective reduction in CLABSI rates.
Study 8	458	Implementation of Enhanced Compliance with Protocols	RR: N/A (Effectiveness: 45%, CI: N/A)	The study concluded a moderately effective in reducing CLABSI.