

Asymptotic Stability of Periodic Solutions for a Nonlinear Neutral First-order Differential Equation with Functional Delay

Amira A. BenFayed

Lecturer, Dept. of Mathematics, Faculty of Education, University of Benghazi, Libya
amira.fayed@uob.edu.ly

Abdelhamid S. Elmabrok

Assistant Professor, Dept. of Mathematics, Faculty of Science, University of
Benghazi, Libya
abdelhamid.elmabrok@uob.edu.ly

Abstract

We consider the nonlinear neutral differential equation:

$$\frac{d}{dt}x(t) = - \prod_{i=1}^p a_i(t) x(t) + \frac{d}{dt} \sum_{i=1}^p Q_i(t, x(t-g(t))) + \int_{-\infty}^t \left[\prod_{i=1}^p D_i(t,s) f(x(s)) + h(s) \right] ds + G(t, x(t), x(t-\tau(t)))$$

and use the contraction mapping principle to show the asymptotic stability of the zero solution provided that $Q(t, 0) = f(0) = 0$.

Keywords: Contraction Mapping, Stability, Nonlinear Neutral, Differential Equation, Integral Equation.

1. Introduction

There have been widely varied solutions for stabilization, Lyapunov's direct methodology being the most renowned. The Lyapunov methodology for the broadly differential equations have been terribly effective in establishing the result for stability see [2,4,5], as well as in establishing the existence of periodic solutions of differential equations with functional delays see [1]. However, there

have been certain issues despite the efficacy of Lyapunov's technique if the functions of equations are unbounded with time and the derivative of the delay is not small and the complexity of generating the Lyapunov function, it is a kind of art to finding this function. Researchers have been working on discovering fresh ways of avoiding those problems. [6] noted that some of these issues disappear when implementing the fixed-point theory. Due to the simplicity of a fixed-point method in comparison with the Lyapunov method, the fixed-point method has become an important instrument to show the existence and uniqueness of solutions and to study the solution's stability in a multitude of mathematical problems. There for, many studies have been published on subject of periodicity and stability of nonlinear neutral equations see [7,8,9,10].

In [3] discussed the existence and uniqueness of periodic solutions of

$$\frac{d}{dt}x(t) = -\prod_{i=1}^p a_i(t)x(t) + \frac{d}{dt}\sum_{i=1}^p Q_i(t, x(t-g(t))) + \int_{-\infty}^t [\prod_{i=1}^p D_i(t,s)f(x(s)) + h(s)] ds + G(t, x(t), x(t-\tau(t))) \quad (1.1)$$

by assuming $a(t)$ is a continuous real-valued function. Taking into consideration

$Q: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $D: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $f: \mathbb{R} \rightarrow \mathbb{R}$, $x: \mathbb{R} \rightarrow \mathbb{R}$, $\mathbf{G}: \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and $h: \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions and to ensure periodicity the following assumption has been made

$a(t)$, $g(t)$, $D(t, x)$, $Q(t, x)$ and $G(t, x, y)$ are periodic functions.

Also, let C_T stand for the set of all continuous scalar functions $x(t)$ periodic in t of period T .

This paper is mainly concerned with the asymptotic stability of the zero solution on the eq.(1.1), as follows we have to mutate eq.(1.1) to an integral mapping equation appropriate for the contraction mapping theorems. This article is organized as follows: Section 2 presents the hypotheses to be used in the later sections, it also introduces Lemma 2.1 that converts eq. (1.1) into an

essential equation, and Section 3 presents the main results.

2. Preliminaries

In this section, we introduce some significant notations. It is appropriate to assume the following conditions. Let $Q(t, x)$, $G(t, x, y)$ and $f(x)$ be globally Lipschitz. So, for $E_1, E_2, E_3, E_4, E_5, E_6$ and K are positive constants such that,

$$\sum_{i=1}^p |Q_i(t, x) - Q_i(t, y)| \leq E_1 \|x - y\| \quad (2.1)$$

$$|f(x) - f(y)| \leq E_2 \|x - y\| \quad (2.2)$$

$$\int_{-\infty}^t \prod_{i=1}^p |D_i(t, s)| ds \leq E_3 < \infty, \quad h(s) \leq E_4 \leq KE_4 \quad (2.3)$$

and,

$$|G(t, x, y) - G(t, w, z)| \leq E_5 \|x - w\| + E_6 \|y - z\| \quad (2.4)$$

The following lemma helps transform eq. (1.1) to an integral corresponding equation.

Lemma 2.1

Let $Q(t, x)$, $D(t, s)$, $a(t)$, $f(t)$, $x(t)$, $g(t)$, $h(t)$ and $G(t, x, y)$ are defined as above, then $x(t)$ is a solution of eq. (1.1) if and only if

$$\begin{aligned} x(t) = & \sum_{i=1}^p Q_i(t, x(t - g(t))) + \left[x(0) - \sum_{i=1}^p Q_i(0, x(0 - g(0))) \right] e^{-\int_0^t \prod_{i=1}^p a_i(k) dk} \\ & + \int_{t-T}^t \left[-\prod_{i=1}^p a_i(u) \sum_{i=1}^p Q_i(u, x(u - g(u))) + \int_{-\infty}^u \left[\prod_{i=1}^p D_i(u, s) f(x(s)) + h(s) \right] ds + \right. \\ & \left. G(u, x(u), x(u - \tau(u))) \right] e^{-\int_0^t \prod_{i=1}^p a_i(k) dk} du \quad (2.5) \end{aligned}$$

Proof.

Let $x(t) \in C_T$ be a solution of eq. (1.1). Now, by writing eq. (1.1) as

$$\frac{d}{dt} x(t) = -\prod_{i=1}^p a_i(t) x(t) + \frac{d}{dt} \sum_{i=1}^p Q_i(t, x(t - g(t))) + \int_{-\infty}^t \left[\prod_{i=1}^p D_i(t, s) f(x(s)) + h(s) \right] ds + G(t, x(t), x(t - \tau(t)))$$

Adding $\prod_{i=1}^p a_i(t) \sum_{i=1}^p Q_i(t, x(t-g(t)))$ to both sides of the last equation, we find:

$$\begin{aligned} \frac{d}{dt} \left[x(t) - \sum_{i=1}^p Q_i(t, x(t-g(t))) \right] &= - \prod_{i=1}^p a_i(t) \left[x(t) - \sum_{i=1}^p Q_i(t, x(t-g(t))) \right] \\ &- \prod_{i=1}^p a_i(t) \sum_{i=1}^p Q_i(t, x(t-g(t))) + \int_{-\infty}^t \left[\prod_{i=1}^p D_i(t, s) f(x(s)) + h(s) \right] ds + \\ &G(t, x(t), x(t-\tau(t))) \end{aligned} \quad (2.6)$$

Now, multiply both sides of (2.6) by $e^{\int_0^t \prod_{i=1}^p a_i(k) dk}$, then integrate from 0 to t, we have:

$$\begin{aligned} &\left[x(t) - \sum_{i=1}^p Q_i(t, x(t-g(t))) \right] e^{\int_0^t \prod_{i=1}^p a_i(k) dk} - \left[x(0) - \sum_{i=1}^p Q_i(0, x(0-g(0))) \right] e^{\int_0^t \prod_{i=1}^p a_i(k) dk} \\ &= \int_0^t \left[- \prod_{i=1}^p a_i(u) \sum_{i=1}^p Q_i(u, x(u-g(u))) + \int_{-\infty}^u \left[\prod_{i=1}^p D_i(u, s) f(x(s)) + h(s) \right] ds + \right. \\ &G(u, x(u), x(u-\tau(u))) \left. \right] e^{\int_0^t \prod_{i=1}^p a_i(k) dk} du \\ &= \int_0^t \left[- \prod_{i=1}^p a_i(u) \sum_{i=1}^p Q_i(u, x(u-g(u))) + \int_{-\infty}^u \left[\prod_{i=1}^p D_i(u, s) f(x(s)) + h(s) \right] ds \right. \\ &\quad \left. + G(u, x(u), x(u-\tau(u))) \right] e^{\int_0^t \prod_{i=1}^p a_i(k) dk} du \end{aligned}$$

Now, by dividing both sides of the above equation by $e^{\int_0^t \prod_{i=1}^p a_i(k) dk}$, we get:

$$\begin{aligned} x(t) &= \sum_{i=1}^p Q_i(t, x(t-g(t))) + \left[x(0) - \sum_{i=1}^p Q_i(0, x(0-g(0))) \right] e^{-\int_0^t \prod_{i=1}^p a_i(k) dk} \\ &+ \int_0^t \left[- \prod_{i=1}^p a_i(u) \sum_{i=1}^p Q_i(u, x(u-g(u))) + \int_{-\infty}^u \left[\prod_{i=1}^p D_i(u, s) f(x(s)) + h(s) \right] ds \right. \\ &\quad \left. + G(u, x(u), x(u-\tau(u))) \right] e^{-\int_0^t \prod_{i=1}^p a_i(k) dk} du \end{aligned}$$

Thus, we see that x is a solution of eq. (1.1).

3. Main Results

The methods employed in this section are adapted from the paper [1] we are supposing that $g(t)$, $f(t)$ and $\tau(t)$ are continuous, $g(t) > 0$ and $Q(t, 0) = f(0) = 0$. Let, $\psi(t): (-\infty, 0] \rightarrow \mathbb{R}$ gives continuous bounded initial function.

We say $x(t) := x(t, 0, \psi)$ is a solution of eq.(1.1) if $x(t) = \psi(t)$ for $t \leq 0$ and satisfies Eq(1.2) for $t \geq 0$.

We say the zero solution of eq.(1.1) is stable at t_0 if for each $\varepsilon > 0$ there is a $\delta = \delta(\varepsilon) > 0$, such that $[\psi : (-\infty, t_0]_T \rightarrow \mathbb{R}$ with $\|\psi\| < \delta$] on $(-\infty, 0]$ implies $|x(t, t_0, \psi)| < \varepsilon$.

$$\int_0^t \prod_{i=1}^p a_i(s) ds > 0 \text{ and } e^{-\int_0^t \prod_{i=1}^p a_i(s) ds} \rightarrow 0, \text{ as } t \rightarrow \infty \quad (3.1)$$

There is an $\alpha > 0$ such that

$$E_1 + \int_0^t [|\prod_{i=1}^p a_i(u)| E_1 + E_2 E_3 + E_4 + E_5 + E_6] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du \leq \alpha < 1 \quad (3.2)$$

$$t - g(t) \rightarrow \infty, \text{ as } t \rightarrow \infty \quad (3.3)$$

$$Q(t, 0) \rightarrow 0, \text{ as } t \rightarrow \infty \quad (3.4)$$

$$G(t, 0, 0) \rightarrow 0, \text{ as } t \rightarrow \infty \quad (3.5)$$

Where, E_1, E_2, E_3, E_4, E_5 and E_6 are defined in inequalities (2.1) _ (2.4).

Theorem 3.1

If the inequalities (2.1) -(2.4) and the conditions (3.1) -(3.5) hold, then every solution $x(t, 0, \psi)$ of eq. (1.1) with small continuous initial function $\psi(t)$ is bounded and approaches zero as $t \rightarrow \infty$.

Moreover, the zero solution is stable at $t_0 = 0$

Proof.

Define the mapping $P: U \rightarrow U$ by $(p\varphi)(t) = \psi(t)$ if $t \leq 0$, and, if $t > 0$.

We have

$$\begin{aligned} (p\varphi)(t) = & \sum_{i=1}^p Q_i(t, \varphi(t - g(t))) + \left[\psi(0) - \sum_{i=1}^p Q_i(0, \psi(0 - g(0))) \right] e^{-\int_0^t \prod_{i=1}^p a_i(k) dk} \\ & + \int_0^t \left[- \prod_{i=1}^p a_i(u) \sum_{i=1}^p Q_i(u, \varphi(u - g(u))) + \int_{-\infty}^u \left[\prod_{i=1}^p D_i(u, s) f(\varphi(s)) + h(s) \right] ds \right. \\ & \left. + G(u, \varphi(u), \varphi(u - \tau(u))) \right] e^{-\int_0^t \prod_{i=1}^p a_i(k) dk} du \end{aligned}$$

It is noticeable that $\varphi \in U$, $(P\varphi)(t)$ is continuous. Let $\varphi \in U$ with $\|\varphi\| \leq K$ for some positive constant K . Let, $\psi(t)$ be small given continuous initial function with $\delta > 0$, $\|\psi\| < \delta$, then using (3.2) in the definition of $(P\varphi)(t)$, we have:

$$\begin{aligned} \|(p\varphi)(t)\| & \leq E_1 k + (1 + E_1) \delta \\ & + \int_0^t \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 K + E_2 E_3 K + E_4 K + (E_5 + E_6) K \right] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du \\ & \leq (1 + E_1) \delta + K \left(E_1 + \int_0^t \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 + E_2 E_3 + E_4 + E_5 + E_6 \right] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du \right) \\ & \leq (1 + E_1) \delta + K \alpha \end{aligned}$$

Which implies that, $\|(P\varphi)(t)\| \leq K$, for the right δ . Thus, (3.4) implies $(P\varphi)(t)$ is bounded. Next, we show that $(P\varphi)(t) \rightarrow 0$ as $t \rightarrow \infty$

The second term on the right side of $(P\varphi)(t)$ tends to zero, by condition (3.1). In addition, the first term on the right side tends to zero, because of (3.3), (3.4) and the fact that $\varphi \in \mathcal{U}$. It is left to show that the integral term goes to zero as $t \rightarrow \infty$

Let $\varepsilon > 0$ be given and $\varphi \in U$ with $\|\varphi\| \leq K$, $K > 0$. Then, there exists a $t_1 > 0$

so that for $t > t_1$, $|\varphi(t - g)(t)| < \varepsilon$. Due to condition (3.1), there exists a $t_2 > t_1$ such that $t > t_2$ implies that $e^{-\int_{t_1}^t \prod_{i=1}^p a_i(k) dk} < \frac{\varepsilon}{\alpha K}$. Thus for $t > t_2$, we have:

$$\begin{aligned} & \left| \int_0^t \left[-\prod_{i=1}^p a_i(u) \sum_{i=1}^p Q_i(u, \varphi(u - g(u))) + \int_{-\infty}^u \left[\prod_{i=1}^p D_i(u, s) f(\varphi(s)) + h(s) \right] ds + \right. \right. \\ & \left. \left. G(u, \varphi(u), \varphi(u - \tau(u))) \right] e^{-\int_0^t \prod_{i=1}^p a_i(k) dk} du \right| \\ & \leq \int_0^{t_1} \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 K + E_2 E_3 K + E_4 K + (E_5 + E_6) K \right] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du \\ & \quad + \int_{t_1}^t \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 \varepsilon + E_2 E_3 \varepsilon + E_4 \varepsilon + (E_5 + E_6) \varepsilon \right] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du \\ & \leq K \int_0^{t_1} \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 + E_2 E_3 + E_4 + E_5 + E_6 \right] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du + \varepsilon \int_{t_1}^t \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 + \right. \\ & \quad \left. E_2 E_3 + E_4 + E_5 + E_6 \right] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du \\ & \leq K e^{-\int_{t_1}^t \prod_{i=1}^p a_i(k) dk} \int_0^{t_1} \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 + E_2 E_3 + E_4 + E_5 + E_6 \right] e^{-\int_u^{t_1} \prod_{i=1}^p a_i(k) dk} du + \alpha \varepsilon \\ & \leq K \alpha e^{-\int_{t_1}^t \prod_{i=1}^p a_i(k) dk} + \alpha \varepsilon \\ & \leq \varepsilon + \alpha \varepsilon \end{aligned}$$

Hence, $(p\varphi)(t) \rightarrow 0$ as $t \rightarrow \infty$.

It remains to show that $(p\varphi)(t)$ is a contraction under the supremum norm.

Theorem 3.2

Let J be a positive constant satisfying the inequality.

$$E_1 + \int_0^t \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 + E_2 E_3 \right] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du \leq J \leq 1 \quad (3.6)$$

then $(P\varphi)(t)$ is a contraction under the supremum norm.

Proof.

$$\begin{aligned} \text{Let } h, \varphi \in \mathcal{U}. \text{ Then } |(ph)(t) - (p\varphi)(t)| & \leq \left\{ E_1 + \int_0^t \left[\left| \prod_{i=1}^p a_i(u) \right| E_1 + \right. \right. \\ & \left. \left. E_2 E_3 \right] e^{-\int_u^t \prod_{i=1}^p a_i(k) dk} du \right\} \|h - \varphi\| \\ & \leq \alpha \|h - \varphi\|. \end{aligned}$$

Therefore, according to the principle of contraction mapping, $(P\varphi)(t)$ is bound

and tends to be zero since t is infinite, moreover, $(P\varphi)(t)$ has a unique fixed point in U that resolves eq. (1.1). The stability of the zero solution at $t_0 = 0$ resulted from simply replacing K with ϵ . That ends the proof.

4. conclusion

In this paper, the nonlinear neutral first-order differential equation with functional delay eq. (1.1) has been transformed into an integral equation by using Lemma 2.1 The integral equation allows us to create a map that enables us to apply the concept of the contraction-map, which ensures us the stability of periodic solutions for a nonlinear neutral first-order differential equation. This allows us to create a map that enables us to apply the concept of the contraction-map, which ensures us the stability of periodic solutions for a nonlinear neutral first-order differential equation with functional delay.

Acknowledgment

The authors would like to express their appreciation to the referees for reading the manuscript, correcting errors, and valuable hints.

References

- [1] Althubiti, S., Makhzoum, H. A. and Raffoul, Y. N. (2013) 'Periodic solution and stability in nonlinear neutral system with infinite delay', *Applied Mathematical Sciences*, 7(133–136), pp. 6749–6764. doi: 10.12988/ams.2013.38462
- [2] Hatvani, L. (1997) 'Annulus arguments in the stability theory for functional differential equations', *Differential and Integral Equations*, 10(5), pp. 975–1002.
- [3] A.A.Ben Fayed, S.M .Alfrgany& H.A. Makhzoum, (2020).The Existence and Uniqueness of Periodic Solutions for Nonlinear Neutral first order differential equation with Functional Delay.*ScienceJournal of Faculty Education*, 2020(8), 211-225.
- [4] Seifert, G. (1973) 'Liapunov-Razumikhin conditions for stability and boundedness of functional differential equations of Volterra type', *Journal of Differential Equations*, 14(3), pp. 424–430. doi: 10.1016/0022-0396(73)90058-2.
- [5] Burton, T.A. (1985) 'Stability and periodic solutions of ordinary and functional-differential equations', *Mathematics in Science and Engineering*, Academic Press, Inc., Orlando, FL, 178.

-
- [6] Burton, T.A. and Furuuchi, T. (2002) ‘Krasnoselskii’s Fixed Point Theorem and Stability’, *Journal of Chemical Information and Modeling*, 53(9), pp. 1689–1699. doi: 10.1017/CBO9781107415324.004.
- [7] H.A .Makhzouma, R.A .Elmansouri, M.A .Bashir . (2020). Stability of Periodic Solutions for nonlinear neutral first_order differential equation with Functional Delay. *Libyan Journal of Basic Science*11(1), 39-47.
- [8] H.A. Makhzouma, A,S.Elmabrok. . (2020). Asymptotic stability of Periodic Solutions for nonlinear neutral first_order differential equation with Functional Delay. *Libyan Journal of Science & Technology*.11:2, 91-93.
- [9] S. Althubiti, H.A. Makhzouma, Y.N. Raffoul, (2013). Periodic Solutions and stability in nonlinear neutral system with infinite delay. *App. Math. Sci* 7(136), 6749-6764.
- [10] A.A.Ben Fayed, R.A .Elmansouri, A.Imhemed, (2023). Stability of Periodic solutions by Krasnoselskii fixed point theorem of neutral nonlinear system of dynamical equation with Variable Delays. *Global Libyan Journal* (67),1-13.